

The Vaginal Microbiota TaqMan™ Assay

Assays
16S
A. vaginae
B. fragilis
BVAB2
C. albicans
C. dubliniensis
C. glabrata
C. krusei
C. lusitaniae
C. parasilosis
C. trachomatis
C. tropicalis
E. coli
E. faecalis
G. vaginalis
H. ducreyi
HSV1
HSV2
L. crispatus
L. gasseri
L. iners
L. jensenii
M. curtisii
M. genitalium
M. hominis
M. mulieris
Megasphera 1
Megasphera 2
N. gonorrhoeae
P. bivia
RnaseP
S. agalactiae
S. aureus
T. pallidum
T. vaginalis
U. urealyticum

The ICD-10 code descriptions on page 1 of this requisition were taken from the official code set issues by the US Department of Health and Human Services, as updated effective October 1, 2020. Depending on the nature of your practice this list may be a useful reference tool; however, it is not a complete list. Please refer to an ICD-10-CM manual for a complete listing. Responsibility for the provision of correct diagnosis code(s) is with the ordering physician. The ordering physician must always determine, for the specific date of service, the appropriate diagnosis code(s) based on the patient's signs and symptoms. ICD-10 codes in effect at the time of service are required by third-party payers to support the medical necessity of the test(s) and/or profile(s) ordered. Physicians are asked to submit all medically appropriate ICD-10 codes when ordering laboratory testing.

Pro-GeneX provides this information only as a customer service; Pro-GeneX does not suggest, and is not meant to suggest, that any of these codes should or should not be used on any specific patient encounter. Pro-GeneX makes no recommendation regarding the use of any particular diagnosis code(s). Diagnosis information should be reflected in the patient's medical record.